

Section 1-1, Mathematics 108

Sets

A set is a collection of objects. There can be uniformity in the collection, eg.

{Orange, Banana, Apple}

or not

{San Francisco, Umbrella, Decimal point}

The items in brackets are called **members** of the set.

In mathematics we are usually interested in sets whose members are mathematical objects.

Examples are:

- Numbers
- Geometric objects, eg. rectangles, triangles
- Polynomial expressions eg. $ax^2 + bx + c$
- Sets, a set can also be a collection of sets

.Set Notation

We can describe a set in a number of ways.

1) Using words: The set of all integers between 1 and 10

2) Using bracket notation to list the members: $\{1,2,3\}$

3) Using bracket notation to describe the set with an expression:

$\{x : x > 5\}$

Read this as "X such that X is greater than 5".

The characters : or | are both used to mean SUCH THAT

$\{x^2 : x \in \mathbb{N}\}$ Read this as "X squared such that X is a member of the set \mathbb{N} .

\mathbb{N} is a symbol we use to mean the Natural numbers.

$$\mathbb{N} = \{1, 2, 3, \dots\}$$

\in is a symbol meaning "is a member of", so

$$1 \in \{1, 2, 3\}$$

Note that here we describe a set which is infinite. The ... is an ellipsis which means continue on indefinitely.

Note: $\{\{1, 2\}, \{3, 4\}\}$ is a set with two members. Each member of the set is a set with two members.

We've already seen that a set can have a finite number of members or an infinite number:

$\{1, 2, 3\}$ - Finite

$\{1, 2, 3, \dots\}$ - Infinite

$\{x : 0 \leq x \leq 1\}$ - Infinite

A set can also have no members. This is called the empty or null set. We can write the null set as either

$\{ \}$ or \emptyset

Note: $\{\emptyset\}$ is not the empty set. It is a set with one member, the empty set.

Operations on Sets

Union

There are a number of standard operations we can perform on sets.

The union of two sets is a set with all the members of the two sets.

$$\{1, 2, 3\} \cup \{3, 4, 5\} = \{1, 2, 3, 4, 5\}$$

The symbol we use looks like a U so it is easy to remember.

Note that the two sets each have the member 3, but it is represented only once in the union. Each member of a set must be unique.

Intersection

The intersection of two sets is a set with only the members that are in both sets.

$$\{1, 2, 3\} \cap \{3, 4, 5\} = \{3\}$$

$$\{1, 2, 3\} \cap \{4, 5\} = \emptyset$$

$$\{1, 2, 3\} \cap \{1, 2, 3\} = \{1, 2, 3\}$$

Sometimes we will represent sets with letters, the same way we use variables in mathematical expressions. Usually we use a capital letter to represent a set, for example:

$$A = \{1, 2, 3\}$$

Using a letter as a variable we can write some simple laws of sets, for example:

$$A = A \cup A$$

$$A = A \cap A$$

Subsets

A subset of a set A is a set B whose members are all in the original set.

The subsets of $A = \{1, 2\}$ are

$$\{ \}, \{1\}, \{2\}, \{1, 2\}$$

Note that the empty set and the set itself are always subsets of a set.

The set and the empty set are called **improper** subsets and all other subsets are called **proper** subsets.

We can write A is a subset of B using the symbols $\subset, \supset, \subseteq, \supseteq$

$$A \subseteq B$$

$$B \supseteq A$$

$$A \subset B$$

$$B \supset A$$

If you think of the \subset as a $<$ you can see which set can have more members.

Sometimes \subset and \supset mean a proper subset.

The set that includes all subsets is called the **power** subset. It can be written this way

$$A = \{1, 2\}$$

$$P(A) = \{ \{ \}, \{1\}, \{2\}, \{1, 2\} \}$$

Note that the power set of a set with 2 elements has 4 elements. Also

$$B = \{1, 2, 3\}$$

$$P(B) = \{ \{ \}, \{1\}, \{2\}, \{3\}, \{1, 2\}, \{1, 3\}, \{2, 3\}, \{1, 2, 3\} \}$$

A set with 3 elements has a power set with 8 elements.

This suggests that a set with n members has 2^n members.

Complement of a Set

The complement of a set A is a set B with all the elements that are not in A.

We write this

$$A^c = B$$

Note that we need to know what elements are being considered. These elements are a set we call U or the **universe** of elements.

$$\text{If } U = \{1, 2, 3, 4\} \text{ and } A = \{1, 2\} \text{ then } A^c = \{3, 4\}$$

Note that if $U=A$ the complement of A is the null set.

Also note that the complement of the null set is always the universe.

Important sets and their symbols

We give these important sets the following symbols:

The natural numbers - $\mathbb{N} = \{1, 2, 3, \dots\}$

The integers - $\mathbb{Z} = \{\dots, -2, -1, 0, 1, 2, \dots\}$ The Z stands for zahlen which is the German word for numbers.

The rational numbers - $\mathbb{Q} = \left\{ \frac{a}{b} : a \in \mathbb{Z}, b \in \mathbb{N} \right\}$ The Q stands for quotient

The real numbers - \mathbb{R}

The complex numbers - $\mathbb{C} = \{a + bi : a, b \in \mathbb{R}, i = \sqrt{-1}\}$

Note that: $\mathbb{N} \subset \mathbb{Z} \subset \mathbb{Q} \subset \mathbb{R} \subset \mathbb{C}$

Properties of the real numbers

We will list some of the properties of addition and multiplication of the real numbers.

Addition

Closure property	if $x, y \in \mathbb{R}$ then $x + y \in \mathbb{R}$
Associate property	if $x, y, z \in \mathbb{R}$ then $(x + y) + z = x + (y + z)$
Commutative property	if $x, y \in \mathbb{R}$ then $x + y = y + x$
Identity property (of 0)	if $x \in \mathbb{R}$ then $x + 0 = 0 + x = x$
Inverse property	if $x \in \mathbb{R}$ then there exists $-x$ such that $x + -x = 0$

Multiplication

Closure property	if $x, y \in \mathbb{R}$ then $xy \in \mathbb{R}$
Associate property	if $x, y, z \in \mathbb{R}$ then $(xy)z = x(yz)$
Commutative property	if $x, y \in \mathbb{R}$ then $xy = yx$
Identity property (of 1)	if $x \in \mathbb{R}$ then $x \cdot 1 = 1 \cdot x = x$
Inverse property	if $x \in \mathbb{R}$ and $x \neq 0$ then there exists $\frac{1}{x}$ such that for all $x \cdot \frac{1}{x} = 1$

Distributive property (of multiplication over addition)

$$x, y, z \in \mathbb{R} \text{ then } x(y + z) = xy + xz$$

Note that subtraction and division do not share all of these properties. Also note that \mathbb{Q} and \mathbb{C} have all of these properties, but \mathbb{N} does not.

Intervals

One important type of subset of the reals is called an **interval**.

An interval is a continuous set of numbers.

You can think of it as a line segment on the real number line.



This describes a **closed** interval $\{x : 1 \leq x \leq 2\}$ which is written $[1,2]$.




This describes a **open** interval $\{x : 1 < x < 2\}$ which is written $(1,2)$.

Note that this is not an ordered pair, or point on an x,y graph.

An interval can also be half open $\{x : 1 \leq x < 2\}$



One end of the interval can be infinite. $\{x : x \geq 1\}$



We write this as $[1,\infty)$. Note that the infinite side is considered open, not closed.

Finally we can write all the real numbers as an interval $\mathbb{R} = (-\infty, \infty)$

Absolute Value

The absolute value is defined as follows

$$|x| = \begin{cases} x & \text{if } x \geq 0 \\ -x & \text{if } x < 0 \end{cases}$$

Some properties of the absolute value

$$|x| \geq 0$$

$$|x| = |-x|$$

$$|x - y| = |y - x|$$

$$|xy| = |x||y|$$

$$\frac{|x|}{|y|} = \frac{|x|}{|y|}$$

These should all be obvious

Finally we have the triangle inequality

$$|x + y| \leq |x| + |y|$$

Note that if x and y are both positive or both negative, you get equality and otherwise you get $<$.

Distance on a line

Let $d(x, y)$ be the distance between two points on the number line, then

$$d(x, y) = |x - y|$$